



Volume XIII

PUP Journal of Science and Technology

January to December 2020

ISSN: 1908-9058

POLYTECHNIC UNIVERSITY OF THE PHILIPPINES

Journal of Science and Technology



Produced by:
Research Institute for Science and Technology

Cover Design by:
Christine Joyce S. Bautista

Layout by:
Jesusana S. Dejito

Technical Assistance by:
Research Publications Office

PUP Journal of Science and Technology

Volume 13, January to December 2020
e-ISSN-2546-0749

Journal Description

The **PUP JOURNAL OF SCIENCE AND TECHNOLOGY (PUPJST)** is a CHED-accredited, double-blind, peer reviewed journal that publishes original articles on theoretical and applied studies in the field of science and technology. It is an annual research publication that aims to provide significant involvement of researchers by presenting novel ideas as well as new knowledge for the advancement of the society in general.

Focus and Scope of Journal

The **PUP JOURNAL OF SCIENCE AND TECHNOLOGY (PUPJST)** publishes original scientific papers and application-focused articles from recent accomplished researches on physical sciences, life sciences, food sciences, nutrition and dietetics, agricultural sciences, environmental sciences, computing and information sciences, mathematics, engineering and technology. The articles must pose valuable hypothesis to derive novel perspectives, verify scientific concepts and innovation in technological advances. The journal does not publish purely descriptive articles that descend from a well-documented exposition, short communication, survey or even a comprehensive review of currently active area of research.

EDITORIAL BOARD MEMBERS

Editor-in-Chief

Armin S. Coronado, *Polytechnic University of the Philippines*

Managing Editor

Alexander S. Carrascal, *Polytechnic University of the Philippines*

Editorial Members/ Advisory Board

Dahlia D.C. Apodaca, *Institute of Human Genetics, National Institutes of Health*

Jerico B. Bacani, *University of the Philippines-Baguio*

Melito A. Baccay, *Technological University of the Philippines*

Lawrence M. Liao, *Hiroshima University, Japan*

Ira C. Valenzuela-Estropia, *De La Salle University*

Associate Editors/ Internal Reviewers

Aleta C. Fabregas, *Polytechnic University of the Philippines*
Alvin R. Dela Cruz, *Polytechnic University of the Philippines*
Bernadeth G. Nobles, *Polytechnic University of the Philippines*
Cristian S. Lazana, *Polytechnic University of the Philippines*
Gary Antonio C. Lirio, *Polytechnic University of the Philippines*
Geoffrey T. Salvador, *Polytechnic University of the Philippines*
Gherhard P. Tan, *Polytechnic University of the Philippines*
Guillermo O. Bernabe, *Polytechnic University of the Philippines*
Ibylou Bandala-Golla, *Polytechnic University of the Philippines*
Jacky Boy E. Atienza, *Polytechnic University of the Philippines*
Jacquiline S. Tychuaco, *Polytechnic University of the Philippines*
John Patrick B. Sta. Maria, *Polytechnic University of the Philippines*
Kevin M. Suliva, *Polytechnic University of the Philippines*
Leana Rich D. Herrera, *Polytechnic University of the Philippines*
Mary Jane M. Tan, *Polytechnic University of the Philippines*
Noel A. Saguil, *Polytechnic University of the Philippines*
Noruel Gerard A. Salvador, *Polytechnic University of the Philippines*
Ria A. Sagum, *Polytechnic University of the Philippines*
Ryan V. Labana, *Polytechnic University of the Philippines*

Layout Artist

Jesusana S. Dejito

External Reviewers/ Referees

Alexa Ray R. Fernando, *National University*
Angel Anne Yanagihara, *University of Hawaii at Manoa*
Argel A. Bandala, *De La Salle University*
Armando Victor M Guidote, *Ateneo de Manila University*
Christian P. Enoval, *Singtel Philippines, Inc.*
Cid Matthew S. Adolfo, *Eulogio "Amang Rodriguez" Institute of Science and Technology*
Cynthia C. Divina, *Central Luzon State University*
Edison A. Roxas, *University of Sto. Tomas*
Gilbert U. Yu, *Ateneo de Manila University*
Jennifer C. Dela Cruz, *Mapua University*
Ma. Vivian D. Camacho, *University of the Philippines-Los Baños*
Michael Y. Roleda, *University of the Philippines-Diliman*
Renan G. Baldovino, *De La Salle University*
Ryan P. Vicerra, *De La Salle University*
Sammy V. Militante, *University of Antique*
Veranoot Nissapatorn, *De La Salle University*

CONTENTS

INFLUENCE OF NITROGEN CONCENTRATION ON THE C-PHYCOCYANIN PRODUCTION OF <i>Spirulina platensis</i> (GOMONT) GEITLER <i>Armin S. Coronado, Florabelle B. Cabbarubias, and Lanieleen Jerah Mae G. Arocha</i>	1
CYTOTOXIC AND APOPTOTIC ACTIVITIES OF MARINE SPONGE <i>Stylissa Massa</i> HEXANE AND METHANOL EXTRACTS AGAINST BREAST CANCER CELL <i>Ramon D. Salanio, Jr., Mary Jho-Anne T. Corpuz, and Ross D. Vasquez</i>	11
STRING EFFICIENCY ANALYSIS OF 132-kV HIGH SUSPENSION INSULATORS USING 2D FINITE ELEMENT METHOD MAGNETICS <i>Federico A. Roy, Jr., Yik Wei Kian, and Alexander S. Carrascal</i>	23
SPECIES LISTING AND SEASONALITY OF MACROFUNGI IN THE CAMPUS OF ISABELA STATE UNIVERSITY, PHILIPPINES <i>James Kennard S. Jacob, Mhark Jelo G. Chavez, Stephanie A. Ignacio, Jose B. Abucay, Jr., and Sofronio P. Kalaw</i>	41
INFLUENCE OF SALINITY IN FATTY ACID PRODUCTION OF <i>Dunaliella</i> sp. AS FEEDSTOCK FOR BIODIESEL <i>John Erasmos Marie P. Talosig, Kia Dyan Louren I. Serrano, and Armin S. Coronado</i>	53

INFLUENCE OF SALINITY IN FATTY ACID PRODUCTION OF *Dunaliella* sp. AS FEEDSTOCK FOR BIODIESEL

JOHN ERASMOS MARIE P. TALOSIG^{1*}, KIA DYAN LOUREN I. SERRANO²,
AND ARMIN S. CORONADO³

¹Philippine Bioresins Corporation

²Department of Education-Disaster Risk Reduction and Management Service

³Director -Research Institute for Science and Technology,
Polytechnic University of the Philippines

*Corresponding Author: ertalosig14@gmail.com

Abstract: *Dunaliella* sp. is a unicellular, eukaryotic, photosynthetic, and halophilic microalga. It is one of the prospective microalgae being utilized for its significant amounts of valuable chemical matters such as carotenoid, glycerol, and lipids. *Dunaliella* has a potential as feedstock in the production of biodiesel but there is only limited data and information available as most studies are focused on commercial production of high-value products such as beta-carotene. Therefore, it is necessary to evaluate the culture condition of *Dunaliella* sp. that promotes higher lipid production. *Dunaliella* sp. was cultivated at varying salinity conditions (45 g/L, 50 g/L, 55 g/L, 60 g/L, 65 g/L, and 70 g/L) using Conway medium (CM). Algal biomass for each culture condition was harvested at the end of logarithmic phase [Period 1 (P1) or day 10] and onset of decline phase [Period 2 (P2) or day 13] of cultivation. At P1, the highest biomass was obtained from CM50 (1.152 ± 0.120 g/L) while CM45 (2.109 ± 0.168 g/L) for P2. Lipid/oil was extracted from algal biomass cultivated at different salinity concentrations by Bligh and Dyer method using solvent mixture of chloroform and methanol (1:2). CM60 has the highest oil yield in both periods, P1 (1.494 ± 0.190 %) and in P2 (1.636 ± 0.173 %). Lipid/Oil from top three (3) oil producing treatments were transesterified to produce the Fatty Acid Methyl Ester (FAME) and subjected to Gas Chromatography - Mass Spectrometry (GC-MS) for fatty acid composition analysis. There are 22 fatty acids identified in P1 while 28 fatty acids in P2. However, only 11 fatty acids identified in all treatments were desirable for biodiesel production namely myristic acid, methyl pentadecanoate acid, palmitic acid, stearic acid, palmitoleic acid, methyl palmitoleate acid, valeric acid, elaidic acid, linoleic acid, oleic acid, and petroselinic acid. Among algal cultures, only CM55 exhibited ideal mix ratio of fatty acid's palmitoleic (16:1), oleic acid (18:1), and myristic acid (14:0) with good biodiesel property (5:3:1). Therefore, *Dunaliella* sp. has a potential as feedstock for biodiesel production with decent amount biomass and lipid/oil using Conway medium that can also exhibit good fuel properties when cultivated at 55g/L of salinity.

Keywords: microalgae, *Dunaliella* sp., lipid, fatty acid methyl ester, biodiesel

1. INTRODUCTION

Globally, an economic growth combined with a rising population has led to the steady increase in energy demands. If the governments around the world stick to current policies, the global energy need will increase to almost 60% in 2030 (IEA, 2007; Patil et al., 2008). Fossil fuels are recognized as the major energy source, but depletion of supply resulted to unsustainable economy (IEA, 2007). Moreover, the continued use of fossil fuels for energy production has contributed to environmental pollution, ecological degradation, and climate change (Hallenbeck & Benemann, 2002; IEA, 2007; Milano et al., 2016). Thus, the modern research community is continuously developing clean and viable energy sources and technologies that are not harmful to the environment (Hallenbeck & Benemann, 2002). Researchers enticed interests to alternative sources for renewable and sustainable energy with high potential for biofuel production such as

microalgae because of higher productivity as compared to land plants as well as its capability to enormously increase its biomass at shorter time (Hu et al., 2008). First generation biofuels are produced from agricultural feedstock that can also be used for food or feed purposes (Mata et al., 2010). The possible competition between food and fuel demand makes it impossible to produce enough biofuel to offset significant percentage of fuel consumption for transportation (Mata et al., 2010; Milano et al., 2016; Peng et al., 2020). Whilst algae appear to be a good candidate as feedstock for biofuel as they can avoid this problem. Also, they lack heavy supporting structures and anchorage organs that can pose technical limitations to their harvesting (Rasoul-Amini, 2014; Peng et al., 2020).

Dunaliella species are green microalgae present in marine environment. Under stress condition, this microalga can accumulate considerable number of valuable metabolites such as carotenoid (Tafreshi & Shariati, 2006), glycerol (Hadi et al., 2008), vitamin and protein (Ghoshal et al., 2002). Moreover, *Dunaliella* sp. offers immense potential in the production of biodiesel, but there is only limited published data and information available as most studies are focused on commercial production of high-value products such as beta-carotene (Tafreshi & Shariati, 2006; Scott et al., 2010; Ahmed et al., 2017). Therefore, it is necessary to evaluate the culture condition of *Dunaliella* sp. that promotes higher lipid production. In this study, *Dunaliella* sp. was cultivated in varying saline concentrations to determine optimal growth condition for biomass and to correlate it with lipid/oil production. Lipid produced by *Dunaliella* sp. cultivated in different saline conditions was characterized by its fatty acid composition profile to provide a reasonable foundation to its potential as biofuel source.

2. METHODOLOGY

2.1 Growth performance of *Dunaliella* sp.

Pure culture of *Dunaliella* sp. was acquired from the Bureau of Fisheries and Aquatic Resources (BFAR) Regional Field Office 1, Alaminos City, Pangasinan and maintained at Phycology Laboratory-Research Institute for Science and Technology (RIST). Prior the experiment, *Dunaliella* was acclimated at different saline concentrations of Conway medium (45g/L to 70 g/L). And then, *Dunaliella* was cultivated at different saline concentrations of Conway medium (CM45=45 g/L, CM50=50 g/L, CM55=55 g/L, CM60=60 g/L, CM65=65 g/L, and CM70=70 g/L) with initial cell density ranges from 1.18×10^5 cells/mL to 1.27×10^5 cells/mL in three (3) replicates. The algal cultures were provided with 24h illumination and continuous aeration. The daily cell density and growth rate were estimated by direct counting using hemocytometer for 13 consecutive days (Martinez-Goss et al., 1975). The total chlorophyll content (TCC) of the cells of each culture was determined on period 1 (P1; Day 10) and period 2 (P2; Day 13) by extracting the chlorophyll using methanol. The absorbance of the extracts was determined using spectrophotometer at wavelength of 650 nm and 665 nm (Martinez-Goss & Dionisio-Sese, 2001).

2.2 Algal wet biomass and lipid extraction

Wet algal biomass from various saline concentrations was harvested by vacuum microfiltration and subsequently centrifuged to obtain the algal pellet. Lipid was

extracted in approximately 2 g of algal pellet using 4.5 mL methanol-chloroform (2:1 v/v). After 2 hours, 1.5 mL of chloroform and deionized water was added, mixed using a vortex mixer, and then centrifuged at 3500 rpm for 10 minutes. The bottom layer was collected and removed the solvent by evaporation to obtain the crude lipid (Bligh & Dyer, 1959).

2.3 Fatty acid profile and characterization

Crude oil extracted from the top three (3) lipid producing cultures was transesterified to produce fatty acid methyl ester (FAME). And then, the fatty acids present were identified by Gas Chromatography Mass Spectrometry (GC-MS) at the National Chemistry Instrumentation Center (NCIC), Ateneo De Manila University. The resulting chromatogram was analyzed and grouped by its chain length (short chain, medium chain, long chain, and very long chain) and degree of saturation (saturated and unsaturated).

2.4 Statistical analyses

All the data gathered for the growth rate, biomass, and lipid production from *Dunaliella* sp. cultivated in varying saline conditions were evaluated using one-way analysis of variance (ANOVA) and Post-hoc Tukey (HSD) using Statistical Program for Social Science (SPSS) version 20 to determine the differences among the treatments. Moreover, correlation between biomass, oil, and salinity concentration was evaluated.

3. RESULTS AND DISCUSSION

3.1 Growth performance of *Dunaliella* sp.

The established growth curve of *Dunaliella* sp. at different salinities is presented in Figure 1. The growth rate (μ^{-1}) of *Dunaliella* sp. cells served as the basis in establishing the phases of cell growth namely lag phase, log or exponential phase, stationary phase, and decline phase. Among the salinity concentrations during log phase (Day 2 - Day 8), CM45 achieved the highest cell density and growth rate at 1.9×10^5 cells/mL and $0.211 \mu^{-1}$, respectively. At stationary phase, CM70 obtained the highest cell density and growth rate at 4.28×10^5 cells/mL and $0.10 \mu^{-1}$, respectively. At the beginning of decline phase, CM60 has the highest cell density and growth rate at 2.84×10^5 cells/mL and $0.054 \mu^{-1}$, respectively.

All the phases of growth are present in all salinity concentrations and observed within the same time frame. *Dunaliella* sp. can survive in a wide range of salinities (Ben-Amotz & Avron, 1992; Tafreshi & Shariati, 2006; Scott et al., 2010; Ahmed et al., 2017). However, rate of cell proliferation and density differs among saline concentrations. The ability of *Dunaliella* sp. to maintain and enhance cell growth varies depending on the salinity. The preferred salinity was higher than 45 g/L or ppt to achieve optimal growth (Abu-Rezq et al., 2010), which is comparable with the result obtained in this study. It was also reported that the optimum salinity condition for *D. salina* and *D. viridis* is 1.0 M NaCl, which is equivalent to 58 g/L (Jimenez & Niell, 1991; Mishra et al., 2008; Hounslow, 2010; Rad et al., 2011). This salinity concentration is within the range of CM

55 and CM 60 used in this study. Moreover, the growth rate of *Dunaliella* sp. is not significantly different among treatments ($p > \alpha_{0.05}$).




Figure 1. Growth curve of *Dunaliella* sp. at different salt concentrations during 13 days of cultivation lag phase (Day 1), exponential phase (Day 2 to Day 8), stationary phase (Day 8 to Day 11), and decline phase (Day 11 to Day 13).

In Figure 2, the summary of the total chlorophyll content (TCC), chlorophyll *a*, and chlorophyll *b* was determined at different salt concentrations (P1 and P2). The highest chlorophyll content was observed during P1, specifically in CM55 ($5.82 \pm 0.87 \mu\text{g/mL}$), followed by CM60 ($5.76 \pm 0.5 \mu\text{g/mL}$) and CM70 ($4.35 \pm 0.17 \mu\text{g/mL}$). Moreover, CM55 also has the highest TCC ($4.88 \pm 0.71 \mu\text{g/mL}$) at period 2 while CM70 has the lowest ($3.8 \pm 0.78 \mu\text{g/mL}$). It is evident that chlorophyll production of *Dunaliella* sp. is inversely related with increasing salinity and with the cultivation period. Additionally, it was observed that chlorophyll *b* is the prominent pigment found in the strain of *Dunaliella* sp. Salinity stress induces change in growth rate, pigmentation, chloroplast structure, and lipid composition in *Dunaliella salina* thus its mortality. It was observed that at higher saline concentration, the decrease in chlorophyll content may be related to chloroplast degradation (Al-Hassan et al., 1987) or low cell growth and proliferation. Salinity is one of the variable factors that changes the yield and composition of algal biomass in *Dunaliella viridis*, due to oxidative stress build-up (Talebi et al., 2013; Hamed et al., 2018). But most of *Dunaliella* species have the capability of adjusting to oxidative stresses by elevating carotene to chlorophyll ratio, which functions as compensatory mechanism. Implying that chlorophyll plays a major role in damage repair processes under stressful conditions (Heraud & Beardall, 2000). This could lead us to the results that chlorophyll content decreased at higher salinity but still capable of proliferating due to its fast adaptation and by altering its chlorophyll composition.




Figure 2. Production of chlorophyll *a*, chlorophyll *b* and total chlorophyll content of *Dunaliella* from different salt concentrations during (A) Period 1 (Day 10) and (B) Period 2 (Day 13).

3.2 Algal wet biomass and lipid production

The summary of total wet algal biomass and percent oil yield for P1 and P2 is presented in Figure 3. Results showed that P2 has the higher values for algal biomass than P1 indicating that proliferation of cells is apparent at P2. In P1, the highest wet biomass was obtained from CM50 (1.152 ± 0.120 g/L). On the other hand, CM45 (2.109 ± 0.168 g/L) got the highest wet biomass in P2. ANOVA revealed no significant differences ($\alpha_{0.05} < 0.942$) among different salinity concentrations in P1. Likewise, there is no significant difference ($\alpha_{0.05} < 0.694$) among different salinity concentrations in P2. This result coincided with the report of Abu-Rezq et al. (2010) that *D. salina* is best grown in 45 ppt. And this study revealed that growth performance of *Dunaliella* sp. is lower in the culture medium with salinity higher than 45 ppt. On the other hand, the highest percent oil yield was obtained from CM65 of P1 (1.494 ± 0.190 %) and P2 (1.636 ± 0.173 %) but there is no significant difference ($\alpha_{0.05} < 0.650$) in P1. Meanwhile in P2, significant difference ($\alpha_{0.05} > 0.030$) between CM60 and CM70 was observed. These results agreed with the findings of Takagi & Karseno (2005) that the optimum starting salinity to produce lipid is around 1.0 M (58 ppt).

Furthermore, there is no significant correlation between wet biomass from P1 ($r = 0.250$, $\alpha_{0.05} < 0.071$) and P2 ($r = 0.135$, $\alpha_{0.05} < 0.335$) with % oil yield from P1 and P2, respectively. CM60 (0.976 ± 0.085 g/L) has the lowest wet biomass among different saline concentrations during P1 but obtained the highest % oil yield (1.494 ± 0.190 %). This implied that *Dunaliella* sp. induces lipid production at high salt concentration although exhibiting lower algal biomass (Takagi & Karseno, 2006; Sonnekus, 2010; Hounslow, 2010). Varying salinity conditions demonstrate the production of lipid from algal cells which is more apparent in moderately high salinities by altering the composition of fatty acids (Sharma et al., 2012; Davis et al., 2015).




Figure 3. Summary of the total wet biomass of *Dunaliella* sp. obtained from various salinity conditions of the culture medium at P1 (10th day) and P2 (13th day) (left chart); Summary and relationship of total wet algal biomass and percent oil yield from *Dunaliella* sp. (A) Period 1 (Day 10) and (B) Period 2 (Day 13) (right chart).

3.3 Fatty acid profile and characterization

3.3.1 Fatty acid composition

CM50, CM55, and CM60 produced the highest lipid content and were used in the characterization of fatty acids (summarized in Table 1). Based on the results, the total number of fatty acids of CM50 decreases from P1 to P2. For CM55, the total SFA increased from 4 SFA in P1 to 7 SFA in P2, along with a decreased in UFA content from 7 UFA in P1 to 5 UFA in P2. For CM60, decreased in total SFA from 7 SFA in P1 to 4 SFA in P2 was observed and an increased in UFA from 16 UFA in P1 to 19 UFA in P2.

Table 1. Fatty Acid Methyl Ester (FAME) profile of *Dunaliella* sp. at P1 and P2 from top producing cultures.

FAME	Common Name	Lipid No.	Period 1			Period 2		
			CM50	CM55	CM60	CM50	CM55	CM60
Saturated								
Methyl tetradecanoate	Myristic acid	C14:0	+	+	+	+	+	-
Pentadecanoic acid, methyl ester	Methyl pentadecanoate	C15:0	+	+	+	+	+	-
Hexadecanoic acid, methyl ester	Palmitic acid	C16:0	+	+	+	+	+	+
Valeric acid, undec-2-enyl ester	Methyl heptadecanoate	C17:0	-	-	+	-	-	-
Methyl stearate	Stearic acid	C18:0	+	+	+	+	+	+
n-Nonadecanol-1	Nonadecanol	C19:0	-	-	-	-	+	-
cis-11-Eicosenoic acid, methyl ester	Eicosenoic acid	C20:0	+	-	+	-	+	-
n-tetracosanol-1	Lignocerosol	C24:0	-	-	-	-	-	+
Tetrahydropyranyl ether of citronellol	Tetrahydropyran	C40:0	+	-	+	-	+	+
Total saturated:			6	4	7	4	7	4
Unsaturated								
Methyl 9-methyltetradecanoate	Methyl tetradecanoate	C14:1 (9)	-	-	-	-	-	+
2-Pentadecanone, 6,10,14-trimethyl-	Pentadecanone	C15:1 (2)	-	-	+	-	-	+
9-Hexadecenoic acid, methyl ester, (Z)-	Palmitoleic acid	C16:2 (9)	+	+	+	+	+	+
Methyl hexadec-9-enoate	Methyl palmitoleate	C16:1 (9)	+	+	+	+	-	+
9-Octadecenoic acid (Z)-, methyl ester	Valeric acid	C16:1 (cis-9)	+	+	+	+	-	-
(R)-(-)-Methyl-8-Hexadecyn-1-ol	Phytol	C16:1	-	-	-	-	+	-
E, E-10,12-Hexadecadien-1-ol acetate	Bombykol	C16:2	-	-	-	-	+	-
7,10-Hexadecadienoic acid, methyl ester	Hexadecadienoic acid	C16:2 (7,10)	-	-	-	-	-	+
4,7,10-Hexadecatrienoic acid, methyl ester	Hexadecatrienoic acid	C16:3	+	+	+	-	-	+
Methyl 4,7,10,13-hexadecatetraenoate	Hexadecatetraenoate	C16:4	-	-	+	+	-	+
cis-11-Eicosenoic acid, methyl ester	Heptadecanol	C17:1	+	-	+	-	-	-
6-Octadecenoic acid methyl ester, (Z)-	Petroselinic acid	C18:1 (6)	-	-	+	-	-	+
9-Octadecen-1-ol, (Z)-	Oleyl alcohol	C18:1 (9-ol)	-	-	-	-	+	-
9-Octadecenoic acid, methyl ester, (E)	Elaidic acid	C18:1(9)	-	-	-	+	+	-
9,12-octadecadienoic acid (Z, Z)-, methyl ester	Linoleic acid	C18:2 (9-12)	+	-	+	+	-	+
n-Heptadecanol-1	Heptadecanol	C18:2	-	+	-	-	-	-
8,11 -octadecadienoic methyl ester	Vaccenic acid	C18:1 (8-11)	-	+	-	-	-	-
8- Octadecenoic acid, methyl ester	Oleic acid	C18:1 (8)	-	+	-	-	-	+
5,8,11,14-Eicosapentaenoic acid, methyl ester	Gondoic acid	C20:1 (5,8,11,14)	+	-	-	-	-	+
(Z)-14-Tricosenyl formate	Tricosenyl	C23:1	-	-	-	-	-	+
5,8,11,14,17-Eicosapentaenoic acid, methyl ester	Omega-3	C25:5	+	-	-	-	-	+
2H-pyran, Tetrahydro-2-(12-pentadecynyloxy)	Narasin	C42:2	+	-	-	-	-	-
11,13-Dihydroxy-tetradec-5-enoic acid,methyl ester	Tetradecenoic acid	N/A	+	-	-	-	-	-
Bicyclo[6.1.0]nonane, 9-(1-methylethylidene)	Nonane	N/A	-	-	-	+	-	+
1H-2-Benzothiopyran, octahydro-, cis-	Benzothiopyran	N/A	-	-	-	-	-	+
Total unsaturated			11	7	9	7	5	15
Total fatty acids:			17	11	16	11	12	19
Total fatty acid per period:			22 Fatty acids			28 fatty acids		

The number of fatty acids in each sample varied from each period and salt concentrations. Result showed that in total, there are 22 fatty acids identified in Period 1 while 28 fatty acids in Period 2. However, only 11 fatty acids are desirable for biodiesel production namely myristic acid, methyl pentadecanoate acid, palmitic acid, stearic acid, palmitoleic acid, methyl palmitoleate acid, valeric acid, elaidic acid, linoleic acid, oleic acid and petroselinic acid. These fatty acids were further clustered into 4 major groups according to its chain length specifically in number of carbon present namely myristic acid group (C14:0), Methyl pentadecanoate group (C15:0), hexadic group (C16: n), and octadic group (C18: n). The relative value to the % oil yield (RV) and group percentage value compared to overall area (GPV) was determined (Table 2). Myristic acid (C14:0), Methyl pentadecanoate, (C15:0) and Hexadic group (C16: n) were relatively high in terms of RV and GPV (Table 2).

Table 2. Summary of computed relative value (RV) and group percentage value (GPV) of top producing cultures from P1 and P2.

Major Groups	Lipid no.	P1			P2			
		CM50	CM55	CM60	CM50	CM55	CM60	
Myristic acid	C14:0	RV	0.072	0.062	0.064	0.054	0.071	-
		GPV	5.06%	4.25%	4.3%	4.23%	4.73%	-
Methyl Pentadecanoate	C15:0	RV	0.014	0.026	0.009	0.014	0.022	0.009
		GPV	0.96%	1.77%	0.58%	1.1%	1.46%	0.59%
Hexadic Group	C16:n	RV	1.03	1.274	0.967	0.913	0.687	1.079
		GPV	72.55%	86.8%	64.76%	71.08%	74.85%	65.94%
Octadic Group	C18:n	RV	0.306	0.017	0.453	0.301	0.285	0.548
		GPV	21.42%	7.18%	30.35%	23.6%	18.97%	33.48%

Likewise, Octadic group (C18: n) is relatively high at CM60 (0.548 and 33.48%) in P2. The result implied that at higher salinity the accumulation of octadic (C18: n) in the cell increases (Zhila et al., 2011). High salinity changes the lipid biosynthetic pathway of the cell and stored in the form of FA and resulted to changes in the fatty acid profile of the cell. Findings in *Botryococcus braunii* and *Nannochloropsis oculata* showed that during high saline stress condition, they undergo elongation of palmitic acid (C16:0) into oleic acid (C18:1), which increases the number of C18 fatty acids in the plasma membrane and promotes desaturation to lessen the permeability of NaCl, preventing the destruction of cells (Kuiper, 1985; Peeler et al., 1989; Fuji et al., 2001; Azachi et al., 2003; Gu et al., 2012). These findings coincided with *Dunaliella* sp., which CM60 got the highest GPV for C18 fatty acids in P1 (30.35%) and in P2 (33.48%). Alongside, CM60 also got the lowest GPV for C16 fatty acids in P1 (64.76%) and P2 (65.94%), respectively.

3.3.2 FAME characteristics

FAME quality of biodiesel is characterized by its chain length and degree of saturation since the composition and amount of FA varied. Hence, it is important to determine the mixture that will promote high quality fuel. It was proposed that the ideal mix ratio of fatty acids at palmitoleic acid (16:1): oleic acid (18:1): myristic acid (14:0) is 5:4:1 (Shenk et al., 2008). This FA ratio gave low oxidative potential, good cold plugging point rating and good cetane number in biodiesel properties. Table 3 presents the computed ratio of FA based on GPV and a close to ideal mix ratio was obtained from CM55 (5:3:1) in P2 which has the same ratio obtained by Fakhry and El Maghraby (2013) from *D. salina*.

Table 3. Summary of computed ratio of fatty acids based on GPV from different salinity in P1 and P2.

	Salinity (ppt)	Fatty Acids		
		C16:1	C18:1	C14:0
P1	CM50	15.99	15.12	5.06
	ratio	3	3	1
	CM55	11.43	6.04	4.25
	ratio	2	1	1
	CM60	15.15	14.73	4.30
	ratio	3	3	1
P2	CM50	17.96	21.49	4.23
	ratio	4	5	1
	CM55	22.50	12.44	4.73
	ratio	5	3	1
	CM60	11.07	14.06	n.a.
	ratio	-	-	-

In P2, CM55 obtained a moderately high GPV of SFA (4.73%) specifically with myristic acid (C14:0) and UFA (22.50%) specifically with palmitoleate acid (C16:1), which described to have good cold plugging point rating and good cetane number. Cold plugging point rating of biodiesel determined by the amount of SFA that has higher melting point than UFA wherein SFA crystallizes at higher temperature compared to UFA. Thus, high SFA promotes higher cold flow and pour point that lessen rapid formation of crystals clogging fuel lines and filters that might cause major operation problems. On the other hand, cetane number is the prime indicator of fuel quality in state-run of diesel engines which is influenced by increasing amount of SFA. Moreover, a low GPV of oleic acid (C18:1) (12.44%) described to have low oxidation. Oxidation stability of biodiesel is determined by the number and position of double bonds in fatty acids. Elevated levels of UFA might result to higher oxidation that will cause fuel to deteriorate. Also, it might pose a critical problem in the storing capacity, but small amount has no reasonable effect on the fuel performance (Lakshminarayanan & Aghav, 2009; Knothe, 2005).

4. CONCLUSIONS

Dunaliella sp. is a green, unicellular, eukaryotic, photosynthetic, and halophilic microalga. It has a potential as feedstock in the production of biodiesel but due to limited published data and information available, development of its commercial production is still dormant. This necessitates the evaluation of the culture condition of *Dunaliella* sp. that promotes higher lipid production with good fuel quality. It is also a highly important undertaking in the development of clean and feasible energy source and technologies that are not harmful to the environment. Thus, the current study determined the optimal

salinity of the culture medium for *Dunaliella* sp. under laboratory condition that produces high yield of lipid with good fuel quality based on its fatty acid composition.

Based on the results, we concluded that *Dunaliella* sp. can grow at different salinities as high as 70 g/L. However, high amount of wet biomass yield was obtained from culture medium with salinity condition of 50 g/L in Period 1 and 45 g/L in Period 2, hence can be considered as the optimum salinity concentrations per period specific for biomass production. On the other hand, total chlorophyll content decreases as salinity concentration increases wherein chlorophyll *b* is more prominent. Chlorophyll plays a major role in damage repair processes under stressful conditions. So, increase in the amount of *chl b* is the probable adaptations of *Dunaliella* sp. to higher saline condition, which can be used in future studies to evaluate the physiological effect of salinity in the growth of *Dunaliella* sp. And then, oil production and fatty acid synthesis of *Dunaliella* sp. varied upon exposure to different saline conditions. The oil production of *Dunaliella* sp. increases as the salinity concentration increases and the optimal salinity concentration specifically for lipid production is 60 g/L. Lastly, 55 g/L salinity (CM55) exhibited good fuel quality using the ideal mix ratio of palmitoleic (16:1), oleic acid (18:1), and myristic acid (14:0) at 5:3:1 ratio. Therefore, *Dunaliella* sp. has a potential as feedstock for biodiesel production with decent amount biomass and lipid/oil using Conway medium that can also exhibit a good fuel property when cultivated at 55g/L of salinity. Research on *Dunaliella* sp. can be enriched by considering other factors that affect the lipid/oil production such as nutrient deprivation, light intensity, and photoperiod. Furthermore, development of new extraction method for higher recovery rate of lipid production is recommended.

5. ACKNOWLEDGEMENT

We would like to express our deepest appreciation to Mr. Gerard C. Khonghun of Salinas Foods, Inc. for funding the materials used for this study. To Antonietta D. Evangelista of BFAR Regional Office 1, Alaminos, Pangasinan, for providing the mother culture used in this study and for the techniques in isolation and cultivation of *Dunaliella* sp. Lastly, to Dr. Elvis Chua of National Chemistry Instrumentation Center, Ateneo de Manila University, for imparting his knowledge, time, and effort with the GC-MS analysis.

6. REFERENCES

- Abu-Rezq, T.S., Al-Hooti, S., & Jacob, D.A. (2010). Optimum culture conditions required for the locally isolated *Dunaliella salina*. *Journal Algal Biomass Utilization*, 1(2), 12-19.
- Ahmed, R. A., He, M., Aftab, R. A., Zheng, S., Nagi, M., Bakri, R., & Wang, C. (2017). Bioenergy application of *Dunaliella salina* SA 134 grown at various salinity levels for lipid production. *Scientific Reports*, 7(1), 1-10.

- Al-Hassan, R. H., Ghannoum, M. A., Sallal, A. K., Abu-Elteen, K. H., & Radwan, S. S., (1987). Correlative changes of growth, pigmentation, and lipid composition of *Dunaliella salina* in response to halostress. *Journal of General Microbiology*, 133(9), 2607-2616.
- Azachi, M., Sadka, A., Fisher, M., Goldshlag, P., Gokhman, I., & Zamir, A. (2003). Salt induction of fatty acid elongase and membrane lipid modifications in the extreme halotolerant alga *Dunaliella salina*. *Plant Physiology*, 132(2), 1115.
- Ben-Amotz, A., & Avron, M. (1992). *Dunaliella: Physiology, biochemistry, and biotechnology*. CRC Press, Boca Raton, p. 240.
- Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8), 911-917.
- Davis, R. W., Carvalho, B. J., Jones, H. D., & Singh, S. (2015). The role of photo-osmotic adaptation in semi-continuous culture and lipid particle release from *Dunaliella viridis*. *J. Appl. Phycol.* 27(1), 109-123.
- Fakhry, E. M., & El Maghraby, D. M. (2013). Fatty acids composition and biodiesel characterization of *Dunaliella salina*. *Journal of Water Resource and Protection*, 5, 894-899.
- Fujii, S., Uenaka, M., Nakayama, S., Yamamoto, R., & Mantani, S. (2001). Effects of sodium chloride on the fatty acids composition in *Boekelovia hooglandii* (Ochromonadales, Chrysophyceae). *Phycological Research*, 49(1), 73-77.
- Ghoshal, D., Mach, D., Agarwal, M., Goyal, A., & Goyal, A. (2002). Osmoregulatory isoform of dihydroxyacetone phosphate reductase from *Dunaliella tertiolecta*: purification and characterization. *Protein Expression and Purification*, 24(3), 404-411.
- Gu, N., Lin, Q., Li, G., Tan, Y., Huang, L., & Lin, J. (2012). Effect of salinity on the growth, biochemical composition, and lipid productivity of *Nannochloropsis oculata* CS 179. *Eng. Life Sci.* 12(5), 631-637.
- Hadi, M. R., Shariati, M., & Afsharzadeh, S. (2008). Microalgal biotechnology: Carotenoid and glycerol production by the green algae *Dunaliella* isolated from the Gave-Khooni salt marsh, Iran. *Biotechnology and Bioprocess Engineering*, 13(5), 540-544.
- Hallenbeck, P. C., & Benemann, J. R. (2002). Biological hydrogen production; fundamentals and limiting processes. *International Journal of Hydrogen Energy*, 27(11-12), 1185-1193.
- Hamed, I., Burcu, A. N., Işın, O., Uslu, L., & Vursavuş, K. K. (2018). The effect of temperature and salinity on the growth and carotenogenesis of three *Dunaliella* species (*Dunaliella* sp. LaNe Isolate, *D. salina* CCAP 19/18, and *D. bardawil* LB 2538) Cultivated under Laboratory Conditions. *Culture*, 19, 18.
- Heraud, P., & Beardall, J. (2000). Changes in chlorophyll fluorescence during exposure of *Dunaliella tertiolecta* to UV radiation indicate a dynamic interaction between damage and repair processes. *Photosynthesis Research*, 63(2), 123-134.
- Hounslow, E. (2010). Optimum salinity conditions for producing lipids from *Dunaliella salina* for biofuels production, Energy Futures Doctoral Training Centre for Interdisciplinary Energy Research. *Mini-project Report*, 5, 13-21.

- Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., & Darzins, A. (2008). Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *The Plant Journal*, 54(4), 621-639.
- International Energy Agency (IEA). 2007. *World Energy Outlook. 2007. China and India Insights*. International Energy Agency Publications: Paris, France.
- Jiménez, C., & Niell, F. X. (1991). Growth of *Dunaliella viridis* Teodoresco: Effect of salinity, temperature and nitrogen concentration. *Journal of Applied Phycology*, 3(4), 319-327.
- Knothe, G. (2005). Dependence of biodiesel fuel properties on the structure of fatty acid alkyl esters. *Fuel Processing Technology*, 86(10), 1059-1070.
- Kuiper, P.J.C. (1985). Environmental changes and lipid metabolism of higher plants. *Physiol. Plant*, 64(1), 118-122.
- Lakshminarayanan, P.A., & Aghav, Y.V. (2009). *Modelling Diesel Combustion*. Springer Science & Business Media.
- Martinez, M.R., Charkoff, R.P., Pantastico, J.B. (1975). Note: Direct phytoplankton counting techniques using the haemocytometer. *Phil. Agri*. 59, 43-60.
- Martinez-Goss, M.R., & Dionisio-Sese, M. L. (2001). Chlorophyll analysis. In: *Analysis of Algae for Food* by Martinez-Goss M.R., Dionisio-Sese, M.L., Migo, V., Elegado, F., Hori, K. (2001) DOST-PCASTRD, Bicutan, Taguig, Metro Manila, Philippines. 27-45.
- Mata, T. M., Martins, A. A., & Caetano, N. S. (2010). Microalgae for biodiesel production and other applications: A review. *Renewable and Sustainable Energy Reviews*, 14(1), 217-232.
- Milano, J., Ong, H. C., Masjuki, H. H., Chong, W. T., Lam, M. K., Loh, P. K., & Vellayan, V. (2016). Microalgae biofuels as an alternative to fossil fuel for power generation. *Renewable and Sustainable Energy Reviews*, 58, 180-197.
- Mishra, A., Mandoli, A., & Jha, B. (2008). Physiological characterization and stress-induced metabolic responses of *Dunaliella salina* isolated from salt pan. *Journal of Industrial Microbiology and Biotechnology*, 35(10), 1093.
- Patil, V., Tran, K. Q., & Giselrød, H. R. (2008). Towards sustainable production of biofuels from microalgae. *International Journal of Molecular Sciences*, 9(7), 1188-1195.
- Peeler, T.C., Stephenson, M. B., Einsphar, K.J., & Thompson Jr, G.A. (1989). Lipid characterization of an enriched plasma membrane fraction of *Dunaliella salina* grown in media of varying salinity. *Plant Physiol*. 89(3), 970-976.
- Peng, L., Fu, D., Chu, H., Wang, Z., & Qi, H. (2020). Biofuel production from microalgae: A review. *Environmental Chemistry Letters*, 18(2), 285-297.
- Rad, F. A., Aksoz, N., & Hejazi, M. A. (2011). Effect of salinity on cell growth and β -carotene production in *Dunaliella* sp. isolates from Urmia Lake in northwest of Iran. *African Journal of Biotechnology*, 10(12), 2282-2289.
- Rasoul-Amini, S., Mousavi, P., Montazeri-Najafabady, N., Mobasher, M. A., Mousavi, S. B., Vosough, F., & Ghasemi, Y. (2014). Biodiesel properties of native strain of *Dunaliella salina*. *International Journal of Renewable Energy Research*, 4(1), 39-41.

- Schenk, P.M., Thomas-Hall, S.R., Stephens, E., Marx, U.C., Mussgnug, J.H., Posten C., Kruse, O., & Hankamer, B. (2008). Second generation biofuels: High-efficiency microalgae for biodiesel production. *Bioenergy Research*, 1(1), 20-43.
- Scott, S. A., Davey, M. P., Dennis, J. S., Horst, I., Howe, C. J., Lea-Smith, D. J., & Smith, A. G. (2010). Biodiesel from algae: Challenges and prospects. *Current Opinion in Biotechnology*, 21(3), 277-286.
- Sharma, K. K., Schuhmann, H., & Schenk, P. M. (2012). High lipid induction in microalgae for biodiesel production. *Energies*, 5(5), 1532-1553.
- Sonnekus, M.J. (2010). *Effects of salinity on the growth and lipid production of ten species of microalgae from the Swartkops Saltworks: A biodiesel perspective* (Doctoral dissertation). Nelson Mandela Metropolitan University, Faculty of Science, p. 98.
- Tafreshi, A. H., & Shariati, M., (2006). Pilot culture of three strains of *Dunaliella salina* for β -carotene production in open ponds in the central region of Iran. *World J. Microbiol. Biotechnol.*, 22(9), 1003–1009.
- Takagi, M., & Karseno, Y. T. (2006). Effect of salt concentration on intracellular accumulation of lipids and triacylglyceride in marine microalgae *Dunaliella* cells. *Journal of Bioscience and Bioengineering*, 101(3), 223-226.
- Talebi, A.F, Tabatabaei, M., Mohtashami, S.K., Tohidfar, M., & Moradi, F. (2013). Comparative salt stress study on intracellular ion concentration in marine and salt-adapted freshwater strains of microalgae. *Not. Sci Biol* 5(3), 309-315.
- Zhila, N.O., Kalacheva, G.S., & Volova, T.G. (2011). Effect of salinity on the biochemical composition of the alga *Botryococcus braunii* Kutz IPPAS H-252. *J. Appl. Phycol.*, 23(1), 47-52.

REVIEW POLICY

An article submitted to the PUP Journal of Science and Technology (PUPJST) undergoes a series of evaluation to ensure adherence to the submission criteria.

1. The Managing Editor pre-screens the article based on its compliance with the SUBMISSION PREPARATION CHECKLIST, and the completeness of the manuscript as stated in the GUIDE FOR AUTHORS. Articles submitted which do not meet all the requirements would be returned to the authors. Upon compliance with the basic requirements, the authors may re-submit the article.
2. PUPJST uses a double-blind peer review system. To maintain its integrity, the author's name and affiliation is deleted in the document. The manuscript shall be assigned a manuscript number that the Section Editors and Reviewers will use as a reference. Section Editors are individuals who are experts in their respective subject areas. The relevant Section Editor completes the article review within 10 working days and may classify the manuscript as:
 - a. accepted as it is;
 - b. accepted with minor revisions;
 - c. accepted with major revisions; or
 - d. rejected
3. Authors shall be informed of the results of the review and are allowed re-submission of the revised article within 15 working days. An article re-submitted beyond the specified time shall already be considered as new submissions. Articles previously rejected by this journal are not accepted for re-submission. The Section Editor recommends the manuscript for evaluation by the Reviewers
4. The recommendations and/or approval of the Reviewer shall be emailed to the author within 15 working days. A manuscript shall be considered for publication upon approval of at least two (2) reviewers.
5. The galley proof of the accepted manuscript will be returned to the author for final proofreading. This galley proof must be returned to the Managing Editor for editing until the submitting author approves a proof.
6. Upon acceptance for publication, copyright of the manuscript is transferred by the author(s) to Polytechnic University of the Philippines. The institution has the exclusive and unlimited rights on the distribution and reproduction of the article.

Submission Preparation Checklist

1. The submission is original and written by the stated author(s).
2. All the authors have read the manuscript and agreed to publish it.

3. The submission has not been previously published nor currently submitted and will not be submitted to another journal while under review by PUPJST.
4. The submission file is in Microsoft Word file format.
5. The text is single-spaced with a 12-point font in Times New Roman and the manuscript text is not in bold face. Each line of the manuscript should be numbered for easy reference during review and corrections.
6. The text employs italics, rather than underlining (except with URL addresses of the references).
7. The text adheres to the stylistic and bibliographic requirements of American Psychological Association (APA). All citations should be reflected in the reference list.
8. Tables, figures, illustrations and other supplementary materials should have a short descriptive title, are properly referenced in the manuscript, and placed with the text or appropriate points rather than at the end.
9. All tables and figures, where applicable, should include all units, clear legends and suitable footnotes.
10. Tables should be in editable format and should not be in graphic objects.
11. Spelling and grammar checks have been performed.
12. After acceptance of the manuscript (during copyediting), proofs in Portable Document File (PDF) format indicating changes and corrections will be sent to the author(s) for approval.

Guide for Authors

1. Only manuscripts that fall within the focus and scope of the Journal will be considered.
2. Manuscripts should be divided into the following sections (in this order):
 - **Title Page**

The title page should provide the title of the article, list the full names and institutional affiliation of all authors and indicate the corresponding author.
 - **Abstract**

The Abstract of the manuscript should not exceed 350 words and must reflect the following parts: background or the context and purpose of the study; results or the main findings; conclusions or brief summary and potential implications. Minimize the use of abbreviations and do not cite references in the abstract.
 - **Keywords**

Three to ten keywords representing the main content of the article
 - **Introduction**

The introduction should be written in a way that is understandable to researchers without specialist knowledge in that area and must clearly state (and if helpful, illustrate) the

background to the research and its aims. The section should end with a brief statement of what is being reported in the article.

Standard chemical symbols and abbreviations may be used in the text, but full term should be given when first mentioned. Units of measurements should be spelled out except when preceded by a numeral. If no-metric measurement units are used the metric equivalent should be mentioned. The complete scientific name of every organism must be cited when it is first mentioned in the text. The generic name may be abbreviated thereafter, except when there are references to other genera with the same initial. The use of common names must be accompanied by the correct scientific name on first use.

- **Methodology**
This section should include the design of the study, the type of materials involved, a clear description of all comparisons, and the type of analysis used, to enable replication. For studies involving human participants, a statement detailing ethical approval and consent should be included.
- **Results and Discussion**
The results and discussion may be combined into a single section or presented separately. This section may also be broken into subsections with short, informative headings.
- **Conclusions**
This should state clearly the main conclusions of the research and give a clear explanation of their importance and relevance. Summary illustrations may be included.
- **Recommendations (if any)**
- **Acknowledgements**
Please acknowledge anyone who contributed towards the article by making substantial contribution to conceptions, design, acquisition of data, or analysis and interpretation of data, or who was involved in drafting the manuscript or revising it critically for important intellectual content, but who does not meet the criteria for authorship. Please also include the source(s) of funding for each author, and for the manuscript preparation. Authors must describe the role of the funding body, if any, in design, in the collection, analysis and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication. Please also acknowledge anyone who contributed materials essential for the study. If a language editor has made significant revision of the manuscript, we recommend that you acknowledge the editor by name, where possible.

Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgements section.

- References

References cited in the text should be presented according to the APA (American Psychological Association) Style Manual, latest edition. The list of References should be given at the end of the paper, immediately following the section on acknowledgement, if any.

3. In preparing illustration and figures, ensure that each figure includes a single illustration and should fit on a single page in portrait format. If a figure consists of separate parts, it is important that a single composite illustration file be submitted which contains all parts of the figure. The following file formats can be accepted: PDF, TIFF, PNG, JPEG or BMP. Note that it is the responsibility of the author(s) to obtain permission from the copyright holder to reproduce figures or tables that have previously been published elsewhere.
4. Proofs will be sent by email to the corresponding author and are expected to proofread the article carefully. The corrected proof should be received by the administration within two working days.
5. The PUPJST adheres to the following four criteria in authorship recommended by International Committee of Medical Journal Editors (ICMJE):
 - a. Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work;
 - b. Drafting the work or revising it critically for important intellectual content;
 - c. Final approval of the version to be published;Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

CONTENTS

e-ISSN-2546-0749 Volume 13

January to December 2020

INFLUENCE OF NITROGEN CONCENTRATION ON THE C-PHYCOCYANIN PRODUCTION OF *Spirulina platensis* (GOMONT) GEITLER

Armin S. Coronado, Florabelle B. Cabbarubias, and
Lanieleen Jerah Mae G. Arocha

1

CYTOTOXIC AND APOPTOTIC ACTIVITIES OF MARINE SPONGE *Stylissa Massa* HEXANE AND METHANOL EXTRACTS AGAINST BREAST CANCER CELL

Ramon D. Salanio, Jr., Mary Jho-Anne T. Corpuz, and
Ross D. Vasquez

11

STRING EFFICIENCY ANALYSIS OF 132-kV HIGH SUSPENSION INSULATORS USING 2D FINITE ELEMENT METHOD MAGNETICS

Federico A. Roy, Jr., Yik Wei Kian, and
Alexander S. Carrascal

23

SPECIES LISTING AND SEASONALITY OF MACROFUNGI IN THE CAMPUS OF ISABELA STATE UNIVERSITY, PHILIPPINES

James Kennard S. Jacob, Mhark Jelo G. Chavez,
Stephanie A. Ignacio, Jose B. Abucay, Jr., and
Sofronio P. Kalaw

41

INFLUENCE OF SALINITY IN FATTY ACID PRODUCTION OF *Dunaliella* sp. AS FEEDSTOCK FOR BIODIESEL

John Erasmos Marie P. Talosig, Kia Dyan Louren I. Serrano, and
Armin S. Coronado

53

